

Decoding postoperative ileus

Miriam Khalil¹ , David G. Jayne¹  and Stephen J. Chapman^{1,*} 

¹Leeds Institute of Medical Research, University of Leeds, Leeds, UK

*Correspondence to: Stephen J. Chapman, St. James's University Hospital, University of Leeds, Room 7.16 Clinical Sciences Building, Leeds LS2 9JT, UK (e-mail: S.Chapman@leeds.ac.uk;  @SJ_Chapman)

Introduction

Postoperative ileus is a temporary cessation of coordinated intestinal motility after surgery. It is characterized clinically as a disruption of normal bowel function in the absence of a mechanical obstruction. It becomes clinically apparent several days after surgery, comprising symptoms of abdominal distension, vomiting, and constipation, often necessitating the placement of a nasogastric tube¹. Postoperative ileus is a common complication, occurring in 10–20% of patients undergoing major abdominal surgery². Patients at greater risk tend to be male, undergoing longer procedures with a higher degree of intestinal handling and surgical stress, and receiving large quantities of opioid-based analgesia after surgery¹. This has a profound impact on recovery after surgery, increasing the risk of venous thromboembolism and hospital-acquired infections. It is also burdensome for healthcare systems, reducing bed capacity and increasing healthcare costs by up to 70%³. Accordingly, the prediction, diagnosis, and treatment of ileus is now considered to be among the highest research priorities in surgery⁴.

Pathogenesis and mechanisms

The physiological and pathological mechanisms leading to postoperative ileus have perplexed healthcare professionals for decades and are still incompletely understood⁵. Broadly, the onset of ileus is considered to comprise two distinct phases: an early neurogenic phase and a later inflammatory phase. During the early phase, stimuli from the initial incision stimulate spinal afferent nerve fibres, activating an inhibitory feedback pathway in the spinal cord which transiently abolishes motility across the intestinal tract. During the inflammatory phase, intestinal handling leads to the release of peritoneal mast cells, increasing intestinal permeability to luminal bacteria and activating intestinal macrophages. Large numbers of nitric oxide- and prostaglandin-releasing leucocytes are then recruited, whose actions impair the contractility of intestinal smooth muscle. Ultimately, this inflammatory cascade leads to the generalized pattern of intestinal dysmotility, characterized clinically as postoperative ileus⁶. The use of opioid analgesia augments these mechanisms by inhibiting propulsive intestinal motility, as do fluctuations in fluid balance, hormonal homeostasis, and electrolyte concentrations⁷.

Perspectives around postoperative ileus

Traditionally, postoperative ileus was considered to be a normal effect of surgery until it was later recognized as a preventable event⁵. Previous research tended to focus on complications considered to be more dangerous for patients and burdensome to health systems, such as anastomotic dehiscence, venous thromboembolism, and surgical site infection. With far greater focus on the patient voice over the last 10 years, however, the true impact of ileus has emerged, illustrated by qualitative studies exploring patients' and health professionals' experiences. Notably, these have highlighted the deleterious effects of ileus on patients, driven physically by its disabling symptoms, such as vomiting and distension, and psychologically through uncertainty related to prolonged recovery. Also highlighted is the disruption to health systems and healthcare professionals, largely due to a lack of proven interventions to reduce ileus and mitigate its effects⁸.

Interventions for postoperative ileus—old, new, and future

Recent advances in perioperative care have generated a range of strategies aimed at reducing the impact and incidence of postoperative ileus (Table 1). Broad evidence supports pathway-based approaches that combine minimally invasive surgery, enhanced recovery after surgery protocols and multimodal analgesia. Evidence for other specific interventions is mixed and often contradictory.

Chewing gum

Chewing gum is inexpensive and generally well tolerated by patients. Enhanced recovery after surgery guidelines provide a moderate recommendation for its use⁹. Its therapeutic potential has been attributed to an efferent cephalic–vagal response and increase in promotility hormones in response to mastication. Numerous studies have explored the clinical effectiveness of chewing gum, with highly variable results. One of the largest studies comprising 1000 patients across 12 hospitals in the Netherlands found that chewing gum did not reduce length of stay or time to intestinal recovery¹⁰.

Probiotics

Probiotics are beneficial bacteria that help to maintain intestinal homeostasis. They exert their effects through various

Received: September 23, 2025. Accepted: September 30, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of BJS Foundation Ltd. All rights reserved. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.



Table 1 Interventions to prevent/treat postoperative ileus

Intervention	Putative mechanism	Typical regimen/practical notes	Evidence/ERAS position & effect
Enhanced Recovery bundle (ERAS core elements: early feeding, multimodal analgesia, minimal drains, early mobilization, fluid optimization)	Multimodal reduction of surgical stress response, earlier gut stimulation and less opioid use	Implement as standard perioperative package (pre → intra → post). Early oral intake as tolerated; multimodal analgesia to minimize opioids; discourage routine NG tubes/drains	Central recommendation of ERAS 2025 consistently associated with reduced POI incidence and shorter time to bowel function
Chewing gum (sham feeding)	Stimulates salivary/gastrointestinal reflexes and vagal tone → promotes bowel motility	10–20 min chew, 3× daily, starting early postoperative day 0–1 until first flatus/bowel motion in many trials	Some RCTs and meta-analyses show shorter time to flatus/bowel movement and shorter LOS; inexpensive and low-risk commonly recommended as adjunct
Early oral/enteral nutrition	Restores luminal stimulation and mucosal integrity; promotes motility	Encourage oral intake as tolerated within 24 h after uncomplicated colorectal surgery	Part of ERAS core; associated with faster return of bowel function and reduced POI
Opioid-sparing/multimodal analgesia (including regional techniques)	Reduces opioid-induced gut hypomotility	Use paracetamol, NSAIDs, gabapentinoids where appropriate, and regional/locoregional blocks to minimize systemic opioids	Strong ERAS recommendation—opioid minimization reduces POI risk
Mu-opioid receptor antagonists (peripheral)—for example alvimopan	Blocks peripheral μ -opioid receptors in gut without reversing central analgesia	Licensed protocol varies by region (preop dose then postoperative dosing for specified days)—availability and cost limit use in some centres	RCTs and meta-analyses show reduced time to GI recovery and LOS after bowel resection; recommended as an option in settings where available
Intravenous lidocaine (systemic)	Anti-inflammatory, antinociceptive and promotility effects when given perioperatively	IV infusion intraop and early postop in selected patients (monitoring required)	Some trials/meta-analyses report faster return of bowel function and pain/opioid reduction; recommended as consideration in appropriate patients
Probiotics/synbiotics	Modulation of gut microbiome and local inflammation → may aid motility recovery	Specific strains/doses vary across studies; usually started peri-/postop	Emerging evidence (some RCTs/meta-analyses) suggests benefit for time to bowel function; evidence heterogeneous—adjunctive option
Minimally invasive surgery (laparoscopic/robotic)	Less surgical trauma and inflammatory response → faster GI recovery	Prefer MIS approaches when appropriate for patient and procedure complexity	Consistently associated with reduced POI and shorter LOS
Avoid routine nasogastric tubes and routine intra-abdominal drains	Prevents iatrogenic delay of enteral intake and reduces discomfort	Use selectively rather than routinely	ERAS advises against routine use after colorectal surgery—reduces factors predisposing to delayed gut recovery
Vagus nerve stimulation/acupuncture/other neuromodulation (experimental)	Augments vagal anti-inflammatory pathway and gut motility	Mostly investigational; some trials use transcutaneous stimulation or acupuncture points	Promising pilot data and biologic plausibility but still experimental—not standard; more high-quality RCTs required
Adjuncts with some supporting data (coffee, dexamethasone, selected herbal formulas, acetylcholinesterase-stimulants in select contexts)	Various—pro-motility, anti-inflammatory, neurohormonal stimulation	Low-cost/low-risk options that have been tested in RCTs/meta-analyses (varied protocols)	Some positive trials for coffee and periop dexamethasone; evidence heterogeneous and generally adjunctive—consider locally and as part of ERAS

Interventions include multimodal perioperative strategies, pharmacological therapies, and adjuncts aimed at reducing surgical stress, opioid-induced gut hypomotility, and inflammation. ERAS® Society Guidelines for Perioperative Care in Elective Colorectal Surgery, 2025 and supporting systematic reviews and meta-analyses. Evidence levels reflect reported outcomes in randomized trials and guideline recommendations. MIS, minimally invasive surgery; LOS, length of stay; NG, nasogastric tube; POI, post-operative ileus.

mechanisms, including competitive inhibition of bacterial adherence and translocation, production of bacteriocins and defensins, and enhancement of barrier function through increased mucus production. Enhanced recovery after surgery

guidelines provide a moderate recommendation for its use⁹. Although high-quality data are lacking, in a meta-analysis of clinical studies ($n=21$), probiotic supplementation was associated with improved intestinal recovery after surgery,

identifying it as a potentially promising intervention to reduce ileus¹¹.

Non-steroidal anti-inflammatory drugs (NSAIDs)

Non-steroidal anti-inflammatory drugs when used as part of multimodal analgesia strategies may reduce the incidence of ileus. This is likely related to a relative reduction in opioid consumption and the attendant impact on intestinal motility. A meta-analysis of six RCTs involving 563 participants previously showed that NSAIDs reduce the time taken for patients to achieve return of bowel function and tolerate an oral diet¹². The use of NSAIDs after colorectal surgery, however, is controversial. Conflicting studies have proposed associations between the use of NSAIDs and anastomotic leak, particularly with non-selective NSAIDs. Enhanced recovery guidelines recommend the use of selective NSAIDs for postoperative analgesia⁹.

Intravenous lidocaine

Local anaesthetic agents are commonly used for epidural analgesia, but recent studies have explored the role of intravenous preparations to reduce ileus. Intravenous lidocaine is thought to attenuate the inflammatory cascade, reduce pain, and decrease postoperative opioid requirements¹. Although several studies have explored its efficacy, most recently the ALLEGRO study showed that perioperative administration of 2% intravenous lidocaine did not improve the return of gut function after minimally invasive colonic resection¹³. Furthermore, the safety of intravenous lidocaine was recently scrutinized, necessitating a consensus statement which recommended a maximum dose, duration, and monitoring precautions when used in the perioperative period¹³.

Mu-opioid antagonists (alvimopam)

Alvimopam is a solution to the adverse effects of opioids on intestinal motility. As a peripherally acting mu-opioid receptor antagonist, alvimopam does not cross the blood-brain barrier and so is considered to mitigate the intestinal effects of opioid without compromising central analgesia. Several high-quality randomized trials have proven the clinical effectiveness of alvimopam for improving intestinal recovery after colorectal surgery^{14,15}. To date, however, the drug is only available in North America.

Vagus nerve stimulation

Vagus nerve stimulation is a novel therapeutic that may reduce ileus through non-invasive stimulation of the cervical or auricular branches of the vagus nerve. Preclinical studies have shown that stimulation of the vagus nerve activates a cholinergic anti-inflammatory pathway, which suppresses intestinal inflammation and prevents smooth muscle dysfunction. Early clinical studies have demonstrated early signals of efficacy, as well as the feasibility of delivering the intervention in clinical practice^{8,16}. Future work is required to explore the clinical and cost-effectiveness of this therapeutic.

Challenges and future directions

Developing and evaluating interventions to reduce ileus is challenging. Specific challenges include community-wide disagreement on the definition of ileus applied to outcome assessment, as well as persisting uncertainty around the mechanism of ileus. This restricts the development of new, evidence-based interventions that have a reasonable chance of

patient benefit. Recent methodological developments, however, may improve our ability to undertake high quality evaluations. Notable examples include a core outcome set for ileus, providing a standard series of patient- and clinician-defined clinical outcomes, as well as a patient-reported outcome measure^{17,18}. Platform trial designs may provide an opportunity to harness this improved methodological framework to facilitate efficient and definitive evaluations of multiple interventions. Before this, however, we must ensure that pipeline early-phase interventions for ileus (such as vagus nerve stimulation) are developed robustly with clear signals of potential efficacy.

Funding

DGJ is supported in part by the National Institute for Health and Care Research (NIHR213331; NIHR205280) and Bowel Cancer UK (18SC0001). SC is supported by the National Institute for Health and Care Research (NIHR CL-2022-02-05) as well as by grants from the The Royal College of Surgeons of England (25PP100030) and the Academy of Medical Sciences (SGL031/1100). The views are those of the authors and not those of the NHS, or Bowel Cancer UK.

Author contributions

Miriam Khalil (Visualization, Writing—original draft, Writing—review & editing), David Jayne (Supervision, Writing—review & editing), and Stephen Chapman (Conceptualization, Supervision, Visualization, Writing—review & editing)

Disclosures

The corresponding author declares: no support from any organisation for the submitted work, no financial relationships with any organizations that might have an interest in the submitted work in the previous three years, and no other relationships or activities that could appear to have influenced the submitted work.

Previous communication

None.

Data availability

All applicable data contained within the manuscript.

References

1. Chapman SJ, Pericleous A, Downey C, Jayne DG. Postoperative ileus following major colorectal surgery. *Br J Surgery* 2018;**105**: 797–810
2. Scarborough JE, Schumacher J, Kent KC, Heise CP, Greenberg CC. Associations of specific postoperative complications with outcomes after elective colon resection: a procedure-targeted approach toward surgical quality improvement. *JAMA Surg* 2017;**152**:e16468
3. Mao H, Milne TGE, O'Grady G, Vather R, Edlin R, Bissett I. Prolonged postoperative ileus significantly increases the cost of inpatient stay for patients undergoing elective colorectal surgery: results of a multivariate analysis of prospective data at a single institution. *Dis Colon Rectum* 2019;**62**:631–637

4. Tiernan J, Cook A, Geh I, George B, Magill L, Northover J *et al.* Use of a modified Delphi approach to develop research priorities for the association of coloproctology of Great Britain and Ireland. *Colorectal Dis* 2014;**16**:965–970
5. Holte K, Kehlet H. Postoperative ileus: a preventable event. *Br J Surgery* 2000;**11**:1480–1493
6. Boeckxstaens GE, de Jonge WJ. Neuroimmune mechanisms in postoperative ileus. *Gut* 2009;**58**:1300–1311
7. Becker G, Blum HE. Novel opioid antagonists for opioid-induced bowel dysfunction and postoperative ileus. *The Lancet* 2009; **9670**:1198–1206
8. Chapman SJ, Kowal M, Helliwell JA, Lockwood S, Naylor M, Croft J *et al.* Non-invasive vagus nerve stimulation to reduce ileus after colorectal surgery: randomised feasibility trial and efficacy assessment (IDEAL stage 2B). *Colorectal Disease* 2024;**12**:2101–2111
9. Gustafsson UO, Rockall TA, Wexner S, How KY, Emile S, Marchuk A *et al.* Guidelines for perioperative care in elective colorectal surgery: Enhanced Recovery After Surgery (ERAS) Society recommendations 2025. *Surgery* 2025;**184**: 109397
10. de Leede EM, van Leersum NJ, Kroon HM, van Weel V, van der Sijp JRM, Bonsing BA *et al.* Multicentre randomized clinical trial of the effect of chewing gum after abdominal surgery. *Br J Surgery* 2018;**7**:820–828
11. Tang G, Huang W, Tao J, Wei Z. Prophylactic effects of probiotics or synbiotics on postoperative ileus after gastrointestinal cancer surgery: a meta-analysis of randomized controlled trials. *PLoS One* 2022;**17**:e0264759
12. Chapman SJ, Garner JJ, Drake TM, Aldaffaa M, Jayne DG. Systematic review and meta-analysis of nonsteroidal anti-inflammatory drugs to improve GI recovery after colorectal surgery. *Dis Colon Rectum* 2019;**2**:248–256
13. Foo I, Macfarlane AJR., Srivastava D, Bhaskar A, Barker H, Knaggs R *et al.* The use of intravenous lidocaine for postoperative pain and recovery: international consensus statement on efficacy and safety. *Anaesthesia* 2021; **76**:238–250
14. Wolff BG, Michelassi F, Gerkin TM, Techner L, Gabriel K, Du W *et al.* Alvimopan, a novel, peripherally acting mu opioid antagonist: results of a multicenter, randomised, double-blind, placebo-controlled, phase III trial of major abdominal surgery and postoperative ileus. *Ann Surg* 2004;**240**: 728–734
15. Delaney CP, Weese JL, Hyman NH, Bauer J, Techner L, Gabriel K *et al.* Phase III trial of alvimopan, a novel, peripherally acting, mu opioid antagonist, for postoperative ileus after major abdominal surgery. *Dis Colon Rectum* 2005;**48**:1114–1125
16. Ru O, Jin X, Qu L, Long D, Liu M, Cheng L *et al.* Low-intensity transcutaneous auricular vagus nerve stimulation reduces postoperative ileus after laparoscopic radical resection of colorectal cancer: a randomized controlled trial. *Minerva Anesthesiol* 2023;**3**:149–156
17. Tripartite Gastrointestinal Recovery Post-operative Ileus Group. Core outcome set for clinical studies of postoperative ileus after intestinal surgery. *Br J Surgery* 2022;**6**:493–496
18. Lee MJ, Baker DM, Hawkins D, Blackwell S, Arnott R, Harji D *et al.* Development of a patient-reported outcome measure for gastrointestinal recovery after surgery (PRO-diGI). *Br J Surgery* 2025;**112**:znaf055